

MIDWEST STONE INSTITUTE

Extracorporeal Shockwave Lithotripsy

In review of the Midwest Stone Institute's ASA/NSAID's list MSI has adopted the following policy with regard to those medications that contain ingredients that may interfere with the blood's ability to clot.

The following medications contain nonsteroidal anti-inflammatory agents. These medications need to be stopped for at least a full 5 days prior to having lithotripsy.

Nonsteroidal Anti-inflammatory Medications:

Advil
Aleve
Anacin
Anaprox
Anaprox DS
Ansaïd
Arthrotec
Cambia
Cataflam
Celebrex
Celecoxib
Clinoril
Co-Advil
Daypro
Diflunisal
Diclofenac
Dolobid
Dristan Sinus Pain Reliever
EC-Naprosyn
Equagesic
Etodolac
Excedrin IB
Excedrin Aspirin Free
Feldene
Fenoprofen
Flector Patch
Flurbiprofen
Genpril

Haltran
Ibuprofen
Ibutab
Indocin
Indomethacin
Ketorolac
Ketoprofen
Lodine
Magsal
Meclofenamate
Meclomen
Medipren
Mefenamic Acid
Meloxicam
Menadol
Midol Extended Relief
Midol Maximum Strength Cramp Formula
Mobic
Motrin
Motrin IB
Nabumetone
Nalfon
Naprelan- Recently Added
Naprosyn
Naproxen
Nuprin
Orudis
Oruvail
Oxaprozin
Pamprin
Piroxicam
Ponstel
Prevacid NapraPAC
Relafen
Rufen
Saletol-200
Sulindac
Tolectin
Tolmetin
Toradol
Trendar
Tricosal
Vioxx
Voltaren
Zipsor

The following medications contain aspirin or salicylic acid derivatives. These medications need to be stopped for at least a full 5 days prior to receiving lithotripsy therapy.

Aspirin or Salicylic Medications:

Aceta-Gesic
Acuprin
Adprin
Aggrenox
Alka Seltzer
Amigesic
Anacin
Anodynos
A.P.C.
Argesic
Artha-G
Arthritis Foundation Pain
Arthritis Pain Formula
Arthritis Strength Bufferin
Arthropan
Arthrotec
ASA
Asacol
Ascriptin
Aspergum
Aspirin
Aspirin with codeine
Asprimox
Azdone
Azulfidine
Backache Maximum
Bayer Products
BC powder and tablets
Bufferin
Bufferin Arthritis Strength
Buffex
Butalbital Compound
Butazolidin
Carna Arthritis
Cataflam
Cogesic
Congesprin

Cope
 Coricidin
 Damason-P
 Darvon Compound
 DeWitt's Pain Reliever
 Dipyridamole
 Disalcid
 Doan's
 Dolene
 Dolor
 Doxaphene
 Dristan
 Durabac
 Duradyne
 Duraxin
 Easprin
 Ecotrin
 Empirin
 Empirin with codeine
 Equagesic
 Equazine-M
 Excedrin
 Extra Strength Anacin
 Femback Caplets
 Fiorina
 Fiortal
 Flexaphen
 Four Way Cold Tablets
 Gelpirin tablets
 Gemnisyn
 Genprin
 Goody's Extra Strength Headache
 Goody's Body Pain
 Halfprin 81
 Heartline
 Halprin
 Healthprin
 Heartline
 Levacet
 Liquiprin
 Lobac
 Lortab ASA
 Magan
 Magnaprin
 Magsal
 Mepor Compound
 Meproamate/Aspirin
 Meprogesic Q
 Marthritic- Recently Added
 Methocarbamol
 Micrainin
 Midol
 Mobidin
 Mobigesic
 Momentum
 Mono-Gesic
 MST 600
 Night-Time Effervescent
 Norgesic
 Norgesic Forte

Norwich Extra Strength
 Nuprin
 Nytol
 Nyquil
 Orphengesic
 Orphenadrine
 Oxycodone and aspirin
 Pabalate-SF
 P-A-C
 Painaid
 Pamprin
 Panasal
 Pentasa
 Pepto-Bismol
 Percodan
 Persantine
 Phenylbutazone
 Phrenillin
 Plavix
 Premysyn
 Presalin
 Quiet World tablets
 Resolve
 Rexolate
 Robaxisal
 Robomol
 Rowasa
 Roxiprin
 S-A-C
 Saletto
 Salflex
 Salocol
 Salsalate
 Salsitab
 Sine-aid
 Sine-off
 Sodium salicylate
 Sodal
 Soma Compound
 Sominex
 St. Joseph Aspirin
 Summit
 Synalgos-DC capsules
 Trental
 Tricosal
 Trigesic
 Trilisate (Choline trisalicylate)
 Tusal
 Vanquish
 Zorprin
The following medications are anticoagulant medications and need to be stopped for the amount of time listed.
Anticoagulant Medications:
(Consult with cardiologist before discontinuing)
COUMADIN (Need PT/PTT) OFF 5 days

WARFARIN (Need PT/PTT)
OFF 5 days
TICLID/TICLOPIDINE (Need Bleeding Time) OFF 14 days
PLAVIX (Check bleeding time) OFF 5 days
EFFIENT (prasugrel) (contact Dr.)
Clopidogrel (contact Dr.)
PLETAL (Need Bleeding time) OFF 5 days
HEPARIN (need PT/PTT) OFF 8 Hours
LOVENOX (enoxaparin) OFF 12 Hours
FRAGMIN (dalteparin) OFF 24 Hours
INNOHEP (tinzaparin) OFF 24 Hours
ARIXTRA (fondaparinux) OFF 5 days (Need Bleeding Time)
REFLUDIN (lepirudin) CHECK PT/PTT
ARGATROBAN CHECK PT/PTT
ANGIOMAX(bivalirudin) (CHECK PT/PTT)
***Arixtra, Refludin, Argatroban, Angiomax : Anticoagulant effect prolonged with renal impairment.**

This list is the most common medications prescribed. There are hundreds of other aspirin containing products on the market that may contain generic aspirin. As such, any medication containing aspirin or acetylsalicylic acid should be stopped for 5 days prior to lithotripsy. If you have any questions regarding these medications, please feel free to call us directly.

HERBAL PRODUCTS AND VITAMINS USE SHOULD STOP BEFORE SURGERY Certain vitamins, minerals and "Herbal medications" can affect anesthetic potency, cause hemodynamic instability, potentiate bleeding, and increase cardiac dysrhythmias. These should be stopped 2 to 3 weeks prior to surgery. (Vitamin E, Echinacea, Ephedra, Feverfew, Garlic, Ginger, Ginko biloba, Ginseng, Goldenseal, Kava-kava, Licorice, Saw Palmetto, St. John's Wort, Valerian).

DO NOT STOP TAKING ANY PRESCRIBED MEDICATIONS WITHOUT CHECKING WITH YOUR PHYSICIAN